


AD A118402

DTIC FILE COPY



TECHNICAL / SCIENTIFIC REPORT

1265

8/82

CEREBROSPINAL FLUID AND SERUM AMPICILLIN LEVELS
IN BACTERIAL MENINGITIS PATIENTS AFTER INTRA-
VENOUS AND INTRAMUSCULAR ADMINISTRATION

By

Isis A. Mikhail, John E. Sippel, Nabil I. Girgis
and Mohamed W. Yassin

DTIC
EL
S
AUG 10 1982
D

U. S. NAVAL MEDICAL RESEARCH UNIT No. 3
(CAIRO, ARAB REPUBLIC OF EGYPT)
FPO NEW YORK 09527

02 00 10 280

**Best
Available
Copy**

Short Communication

Cerebrospinal Fluid and Serum Ampicillin Levels in Bacterial Meningitis Patients after Intravenous and Intramuscular Administration*

ISIS A. MIKHAIL, JOHN E. SIPPEL,
NABIL I. GIRGIS and MOHAMED W. YASSIN

From the US Naval Medical Research Unit No. 3, and Abbassia Fever Hospital, Cairo, Egypt

ABSTRACT. 28 patients with bacterial meningitis received ampicillin by the intramuscular (IM) route and 16 patients by the intravenous (IV) route. The mean cerebrospinal fluid (CSF) ampicillin levels were similar in the two groups 1 h after a dose given on the first or second day of treatment, but they were higher in the IM group on both days 4 h after a dose. CSF/serum ratios were similar in both groups but considerably higher at 4 h than at 1 h.

INTRODUCTION

Ampicillin has been shown to reach therapeutically significant levels in cerebrospinal fluid (CSF) (5) and to be effective in the treatment of bacterial meningitis (4, 5) when it is administered intravenously (IV). However, intramuscular (IM) administration would simplify management of patients, especially in areas where modern medical facilities are not readily available. Since little has been reported on the CSF levels that can be obtained after IM administration in patients with inflamed meninges, ampicillin concentration in blood and CSF specimens from patients with bacterial meningitis receiving IM or IV ampicillin during the first 2 days of therapy were compared.

PATIENTS AND METHODS

44 patients (Table I), whose admission CSF and serum samples contained no detectable antibiotics, were given 160 mg/kg/day of ampicillin divided into 6 doses at 4-h intervals. The first 10 patients were given the drug by the IM route, after which IM or IV administration was used according to a predesigned randomization table. 28 patients received the drug by the IM route for 5-7 days and 16 patients by the IV route for 2 days followed by IM for 5 days. Both groups were then given oral ampicillin for 1 week. CSF and sera were obtained from 16 patients: 11 patients 1 h after the first dose, 13 patients 4 h after the first dose, 14 patients 1 h after the 7th dose, and 17 patients 4 h after the 7th dose. 16 patients had CSF and sera obtained on both day 1 and day 2 (Table II).

The simplified agar-well diffusion method (2) using *Ba-*

cillus subtilis spore suspension (Difco Laboratories) was employed in the determination of ampicillin concentrations in the CSF and sera with simultaneous parallel comparison to antibiotic standards ranging from 0.1 to 80.0 µg/ml. The standards and specimens were stored together at -70°C until tested. Each assay involved 3 replicates of the test specimens and standards.

RESULTS

The range and mean CSF and serum ampicillin levels are given in Table II. The mean CSF values obtained 1 h after IM or IV administration were comparable on days 1 and 2. However, the mean CSF levels 4 h after administration were significantly higher in the patients who received the drug by the IM route both on day 1 ($p < 0.02$) and day 2 ($p < 0.01$). The CSF/serum ratios ($\times 100$) were approximately 5 one hour after ampicillin administration by either route on both days. These ratios rose to approximately 50 at 4 h after administration.

There was no noticeable difference in the clinical response with IM or IV administration. One patient with meningococcal meningitis and one patient with meningitis of unknown etiology died in the IM group, and one patient with meningococcal meningitis died in the IV group.

*The opinions and assertions contained herein are the private ones of the authors and are not to be construed as official or as reflecting the views of the Department of the Navy or the naval service at large.

Table I. Bacterial meningitis patients included in ampicillin study

	Intramuscular	Intravenous
Etiology		
Neisseria meningitidis	18	9
Streptococcus pneumoniae	4	2
Haemophilus influenzae	1	
Staphylococcus aureus		1 ^a
Unknown	5	4
Age (range)	5 mo - 20 yr	4 mo - 62 yr
Age (mean)	7.7 yr	10.5 yr
Male/female ratio	1.3	0.7

^a Patient with Staph. aureus meningitis was given staphicillin IM after 2 days of ampicillin therapy

DISCUSSION

Little data are available on the ampicillin CSF levels achieved after IM administration. Barrett et al. (1), had mean concentrations of 0.43 µg/ml in normal subjects and 0.65 µg/ml in patients with viral meningitis. Thrupp et al. (6) gave ampicillin IM in patients with purulent meningitis but only after they responded satisfactorily with IV therapy. In patients with *haemophilus meningitis*, Wilson and Haltalin (7) demonstrated mean CSF ampicillin levels of 2.0 µg/ml 1 h after an IV dose, 2.1 µg/ml 4 h after an IV dose, 4.7 µg/ml 1 h after an IM dose, and 4.1 µg/ml 4 h after an IM dose during the second day of ampicillin therapy. Their data are consistent with this report, showing that the CSF levels are higher 4 h after IM administration.

IM administration of ampicillin resulted in concentrations higher than the MICs for the usual agents of bacterial meningitis (3, 6; Mikhail et al., unpublished data). IM ampicillin therefore appears to be an effective therapeutic regimen for bacterial meningitis and may be particularly suitable for outbreaks of meningococcal disease in developing countries.

ACKNOWLEDGEMENTS

We thank Dr Michael Kilpatrick for his helpful suggestions regarding this manuscript and Miss Alexandra Patsalidis for its preparation.

This work was supported by the Naval Medical Research and Development Command, Bethesda, Maryland, under Work Unit No. M0095. PN002-5069.

Scand J Infect Dis 13

Table II. Cerebrospinal fluid and serum ampicillin levels in bacterial meningitis patients

	No. of pat.	CSF levels ($\mu\text{g/ml}$)		Serum levels ($\mu\text{g/ml}$)	
		Range	Mean	Range	Mean
Intramuscular route					
Day 1					
1 h	10	0.10-1.65	1.06	10.75-30.50	21.42
4 h	9	0.44-9.50	3.64	<0.10-33.00	8.54
Day 2					
1 h	9	0.23-1.95	1.02	8.00-35.00	19.50
4 h	11	0.42-8.40	2.94	1.60-20.50	7.27
Intravenous route					
Day 1					
1 h	6	0.31-2.50	1.09	10.00-28.50	17.83
4 h	4	0.22-2.90	1.02	0.30-6.00	2.30
Day 2					
1 h	5	0.52-0.82	0.66	1.85-27.00	13.37
4 h	6	0.20-1.70	0.67	0.40-4.50	1.70

REFERENCES

- Barrett, F. F., Eardley, W. A., Yow, M. D. & Leve-rett, H. A.: Ampicillin in the treatment of acute suppurative meningitis. *J Pediatr* 69: 343, 1966.
- Bennett, J. V., Brodie, J. L., Benner, E. J. & Kirby, W. M. M.: Simplified accurate method for antibiotic assay of clinical specimens. *Appl Microbiol* 14: 170, 1966.
- Devine, L. F. & Hagerman, C. R.: Spectra of susceptibility of *Neisseria meningitidis* to antimicrobial agents in vitro. *Appl Microbiol* 19: 329, 1970.
- Girgis, N. I., Yassin, M. W., Sanborn, W. R., Burdick, K. L., El Ella, H. A., Kent, D. C., Sorensen, K. & Nabil, I. M.: Ampicillin compared with penicillin and chloramphenicol combined in the treatment of bacterial meningitis. *J Trop Med Hyg* 75: 154, 1972.
- Hassan, A., Yassin, M. W., Girgis, N. I., Sippel, J. E., Mmer, W. F., Tadros, S. S. & El Ella, H. A.: The value of cepicillin and ampicillin in the treatment of meningococcal and pneumococcal meningitis. *Scand J Infect Dis* 8: 181, 1976.
- Thrupp, L. D., Leedom, J. M., Ivler, D., Wehrle, P. F., Portnoy, B. & Mathies, A.: Ampicillin levels in the cerebrospinal fluid during treatment of bacterial meningitis. In: *Antimicrob Agents Chemother.* p. 206. American Society for Microbiology, Washington 1966.
- Wilson, H. D. & Haltalin, K. C.: Ampicillin in *Haemophilus influenzae* meningitis. Clinico-pharmacologic evaluation of intramuscular vs intravenous administration. *Am J Dis Child* 129: 209, 1975.

Address for reprints:

Research Publications,

NAMRU-3,

FPO New York 09527/0007.


USA

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER NAMRU-3-TR-8/82	2. GOVT ACCESSION NO. A118402	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) CEREBROSPINAL FLUID AND SERUM AMPICILLIN LEVELS IN BACTERIAL MENINGITIS PATIENTS AFTER INTRA- VENOUS AND INTRAMUSCULAR ADMINISTRATION		5. TYPE OF REPORT & PERIOD COVERED Technical/Scientific Report
		6. PERFORMING ORG. REPORT NUMBER Acc. 1265
7. AUTHOR(s) Isis A. Mikhail, John E. Sippel, Nabil I. Girgis and Mohamed W. Yassin		8. CONTRACT OR GRANT NUMBER(s)
9. PERFORMING ORGANIZATION NAME AND ADDRESS U.S. Naval Medical Research Unit No.3 FPO New York 09527		10. PROGRAM ELEMENT PROJECT, TASK AREA & WORK UNIT NUMBERS M0095.PN002-5069
11. CONTROLLING OFFICE NAME AND ADDRESS Naval Medical Research and Development Command National Naval Medical Center Bethesda, MD 20814		12. REPORT DATE 1981
		13. NUMBER OF PAGES 2
14. MONITORING AGENCY NAME & ADDRESS (If different from Controlling Office)		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Distribution of this report is unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES Published in : Scand. J. Infect. Dis. 13:237-238, 1981		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Bacterial Meningitis Cerebrospinal fluid (CSF) CSF/serum ratio		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) 28 patients with bacterial meningitis received ampicillin by the intra- muscular (IM) route and 16 patients by the intravenous (IV) route. The mean cerebrospinal fluid (CSF) ampicillin levels were similar in the two groups 1 h after a dose given on the first or second day of treatment, but they were higher in the IM group on both days 4 h after a dose. CSF/serum ratios were similar in both groups but considerably higher at 4 h than at 1 h.		

Accession	
Indexing	
Classification	
Availability Codes	
Avail. and/or	
Dist. Special	
A 20	



DD FORM 1473

1 JAN 73


EDITION OF 1 NOV 65 IS OBSOLETE

S/N 0102-LF-014-6601

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)



S/N 0102- LF-014-6601

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)